

**Public Health District Health Indicators
2007 Data**

Item	Idaho	District 1	District 2	District 3	District 4	District 5	District 6	District 7
Maternal/Child								
1) % unintended pregnancies	36.3	38.0	34.7	42.1	36.2	38.1	37.9	27.4
2a) % live births with adequate prenatal care in 2007 ¹	42.6	42.6	49.2	39.5	47.9	42.7	41.6	36.3
2b) % live births with intensive prenatal care in 2007 ¹	30.1	35.2	24.3	26.5	30.4	27.5	35.5	30.6
2c) % live births with adequate or intensive prenatal care in 2007 ¹	72.7	77.8	73.4	66.0	78.3	70.2	77.1	66.8
3a) % live births with adequate prenatal care in 2006 ¹	42.0	46.6	46.6	36.3	43.0	46.1	39.5	40.8
3b) % live births with intensive prenatal care in 2006 ¹	31.9	34.9	25.4	29.6	35.0	28.4	38.4	27.7
3c) % live births with adequate or intensive prenatal care in 2006 ¹	73.9	81.5	72.0	65.9	78.0	74.5	77.8	68.4
4) % live births with low birth weight ²	6.6	6.0	6.0	5.8	6.9	6.9	7.3	6.7
5) % live births with tobacco use during pregnancy ³	12.0	21.6	15.6	10.7	12.0	12.8	11.5	6.3
6) % WIC participation ⁴	37.0	37.4	37.2	42.2	23.5	42.1	50.0	40.4
7) % prevalence of breastfeeding ⁵	88.1	88.3	89.8	88.2	89.8	85.0	83.6	89.7
Adolescents								
1) Teen pregnancy rate (15-19) ⁶	49.5	44.8	29.4	75.8	47.4	67.3	45.4	33.2
2) Motor vehicle crash death rate, 2005-2007 (ages 15-19) ^{7,9}	27.7	18.4	35.0	41.1	5.0	54.0	40.5	24.2
3) Suicide rate, 2005-2007 (ages 10-18) ^{8,9}	6.4	10.0	8.3	6.0	3.3	8.4	7.0	6.1

Item	Idaho	District 1	District 2	District 3	District 4	District 5	District 6	District 7
Adults								
1) % without health care coverage	18.9	18.4	15.5	24.1	15.9	24.7	18.4	16.7
2) % with no leisure time activity	19.6	18.1	20.7	23.2	16.4	24.4	21.5	16.9
3) % overweight (BMI ≥ 25)	63.1	60.8	64.6	67.1	64.2	62.9	61.7	58.4
4) % diagnosed with diabetes	7.9	7.0	8.2	9.3	7.2	10.0	8.3	6.3
5) % who smoke	19.1	23.6	19.4	20.6	17.5	23.0	19.1	11.5
6) % who binge drink ¹⁰	14.7	17.4	15.8	13.4	16.8	13.5	15.0	8.7
7) % of females without breast cancer screening	n/a	n/a	n/a	n/a	n/a	n/a	n/a	n/a
8) % of males without prostate cancer screening	n/a	n/a	n/a	n/a	n/a	n/a	n/a	n/a
9) % who did not wear seatbelts	24.6	22.2	23.6	22.3	20.7	25.7	32.8	31.5
10) Suicide rate, 2005-2007 (ages 65+) ^{8,9}	21.1	28.0	32.2	15.9	18.0	21.9	11.0	24.1
11) % with no dental visit in the past 12 months	32.3	36.7	41.0	32.7	25.3	34.1	35.3	32.9

"na" indicates data are not available for this year.

SOURCES:

For Maternal/Child indicator 1: Pregnancy Risk Assessment Tracking Systems (PRATS), Bureau of Vital Records and Health Statistics. PRATS is a representative sample of resident women aged 18+ who gave birth in Idaho.

For Maternal/Child indicators 2-7, Adolescents indicators, and Adults indicator 10: Vital Statistics, Bureau of Vital Records and Health Statistics.

For Adults indicators 1-9, 11: Idaho Behavioral Risk Factor Surveillance System (BRFSS), Bureau of Vital Records and Health Statistics.

NOTES:

1. In 2004, the Idaho birth certificate was revised. Beginning in 2004, Idaho prenatal care data are based on date of first prenatal care visit as reported in the mother's medical record. Prenatal care data in 2004 and after are not comparable with prenatal care based on month prenatal care began. The APNCU index classifies care as intensive, adequate, intermediate, inadequate, or no care by comparing the number of actual prenatal care visits that a woman was expected to receive, given the onset of prenatal care and the length of gestation. Percents based on records with stated data for APNCU calculation (date of first prenatal care visit, number of visits, and length of gestation).
2. Low birth weight: live birth weighing under 2,500 grams (5 1/2 pounds or less). Percents based on records with stated birth weight.
3. Prior to 2004, tobacco use during pregnancy (yes or no) was reported on the Idaho birth certificate. Beginning in 2004, cigarette smoking before and during pregnancy is reported on the revised birth certificate. Data prior to 2004 are not comparable with data in 2004 and after.
4. The revised 2004 Idaho birth certificate includes a new variable for reporting if mother received WIC food for herself during this pregnancy (yes or no)?
5. The revised 2004 birth certificate includes a new variable for reporting if infant was being breastfed at birth or at the time the birth certificate was completed (yes or no)?
6. Pregnancies include live births, induced abortions, and reportable stillbirths (only those fetal deaths with a gestational period of 20+ weeks or which weight 350+ grams are required to be reportable under Idaho law.) Rate: Number of pregnancies per 1,000 females in corresponding age group. Rates are based on July 1, 2007 female population estimates from the National Center for Health Statistics, United States resident population from the Vintage 2007 postcensal series, prepared under a collaborative arrangement with the U.S. Census Bureau, Internet release date September 5, 2008.
7. Motor Vehicle Crash Death Rate: number of motor vehicle crash deaths from 2005 to 2007 in age group per 100,000 population in age group.
8. Suicide rate: number of suicides from 2005 to 2007 in age group per 100,000 population in age group.
9. Rates are based on the National Center for Health Statistics, estimates of the July 1, 2006 United States resident population from the Vintage 2006 postcensal series, prepared under a collaborative arrangement with the U.S. Census Bureau, Internet release date August 16, 2007.
10. Prior to 2005, Idaho BRFSS reported drinking status among all adults (binge drinking = 5 or more drinks in single occasion). Beginning in 2005, Idaho reports gender specific status (5 or more for males, 4 or more for females, in single occasion).

**Public Health District Health Indicators
2006 Data**

Item	Idaho	District 1	District 2	District 3	District 4	District 5	District 6	District 7
Maternal/Child								
1) % unintended pregnancies	34.4	39.3	35.3	38.6	30.7	38.3	37.3	28.3
2a) % live births with adequate prenatal care in 2006 ¹	42.0	46.6	46.6	36.3	43.0	46.1	39.5	40.8
2b) % live births with intensive prenatal care in 2006 ¹	31.9	34.9	25.4	29.6	35.0	28.4	38.4	27.7
2c) % live births with adequate or intensive prenatal care in 2006 ¹	73.9	81.5	72.0	65.9	78.0	74.5	77.8	68.4
3a) % live births with adequate prenatal care in 2005 ¹	41.9	44.4	47.5	36.3	42.5	45.8	39.1	42.5
3b) % live births with intensive prenatal care in 2005 ¹	32.3	34.9	30.5	29.0	35.6	28.0	38.2	27.7
3c) % live births with adequate or intensive prenatal care in 2005 ¹	74.2	79.3	78.0	65.3	78.1	73.8	77.3	70.2
4) % live births with low birth weight ²	6.9	5.6	5.5	7.1	7.6	7.0	7.1	6.7
5) % live births with tobacco use during pregnancy ³	12.7	21.6	18.1	12.2	12.5	12.8	12.4	6.2
6) % WIC participation ⁴	35.9	38.8	39.9	41.3	19.4	41.1	50.5	40.7
7) % prevalence of breastfeeding ⁵	88.4	89.0	89.7	89.2	89.8	85.1	83.4	90.4
Adolescents								
1) Teen pregnancy rate (15-19) ⁶	46.5	42.4	32.9	75.0	46.3	58.3	39.4	28.6
2) Motor vehicle crash death rate, 2004-2006 (ages 15-19) ^{7,9}	27.0	25.3	32.3	27.7	7.6	55.7	44.7	19.3
3) Suicide rate, 2004-2006 (ages 10-18) ^{8,9}	5.8	5.2	11.5	6.3	4.1	7.2	4.3	6.3

Item	Idaho	District 1	District 2	District 3	District 4	District 5	District 6	District 7
Adults								
1) % without health care coverage	18.8	20.6	21.8	23.5	16.3	24.2	13.2	14.0
2) % with no leisure time activity	20.8	16.8	22.2	25.0	17.3	25.7	21.8	21.4
3) % overweight (BMI ≥ 25)	59.7	62.7	61.7	61.1	57.4	58.9	61.7	57.5
4) % diagnosed with diabetes	6.8	7.0	6.8	7.9	6.6	5.5	6.9	7.0
5) % who smoke	16.8	19.2	21.8	20.9	15.3	18.3	15.0	9.3
6) % who binge drink ¹⁰	14.8	17.4	19.6	10.4	18.0	18.2	10.4	7.5
7) % of females without breast cancer screening	37.8	38.2	36.0	39.5	32.7	40.9	43.3	39.4
8) % of males without prostate cancer screening	48.7	49.6	53.0	46.8	48.7	49.7	45.5	48.2
9) % who did not wear seatbelts	26.8	22.2	25.9	23.9	19.7	34.8	37.1	35.9
10) Suicide rate, 2004-2006 (ages 65+) ^{8,9}	22.6	28.8	37.7	22.9	16.2	23.9	11.1	22.7
11) % with no dental visit in the past 12 months	34.6	37.8	39.6	41.7	27.1	43.2	32.4	29.0

"na" indicates data are not available for this year.

SOURCES:

For Maternal/Child indicator 1: Pregnancy Risk Assessment Tracking Systems (PRATS), Bureau of Vital Records and Health Statistics. PRATS is a representative sample of resident women aged 18+ who gave birth in Idaho.

For Maternal/Child indicators 2-7, Adolescents indicators, and Adults indicator 10: Vital Statistics, Bureau of Vital Records and Health Statistics.

For Adults indicators 1-9, 11: Idaho Behavioral Risk Factor Surveillance System (BRFSS), Bureau of Vital Records and Health Statistics.

NOTES:

1. In 2004, the Idaho birth certificate was revised. Beginning in 2004, Idaho prenatal care data are based on date of first prenatal care visit as reported in the mother's medical record. 2004 and 2005 APNCU data are not comparable with APNCU data based on month prenatal care began. The APNCU index classifies care as intensive, adequate, intermediate, inadequate, or no care by comparing the number of actual prenatal care visits that a woman was expected to receive, given the onset of prenatal care and the length of gestation. Percents based on records with stated data for APNCU calculation (date of first prenatal care visit, number of visits, and length of gestation). Data for 2004 have been corrected from the 2004 Health Indicators; 2004 data are comparable to 2005 data.
2. Percents based on records with stated birth weight.
3. Prior to 2004, tobacco use during pregnancy (yes or no) was reported on the Idaho birth certificate. Beginning in 2004, cigarette smoking before and during pregnancy is reported on the revised birth certificate. Data in 2004 and 2005 are not comparable to data prior to 2004.
4. The revised 2004 Idaho birth certificate includes a new variable for reporting if mother received WIC food for herself during this pregnancy (yes or no)?
5. The revised 2004 birth certificate includes a new variable for reporting if infant was being breastfed at birth or at the time the birth certificate was completed (yes or no)?
6. Pregnancies include live births, induced abortions, and reportable stillbirths (only those fetal deaths with a gestational period of 20+ weeks or which weight 350+ grams are required to be reportable under Idaho law.) Rate: Number of pregnancies per 1,000 females in corresponding age group. Rates are based on July 1, 2006 female population estimates from the National Center for Health Statistics, United States resident population from the Vintage 2006 postcensal series, prepared under a collaborative arrangement with the U.S. Census Bureau, Internet release date August 16, 2007.
7. Motor Vehicle Crash Death Rate: number of motor vehicle crash deaths from 2004 to 2006 in age group per 100,000 population in age group.
8. Suicide rate: number of suicides from 2004 to 2006 in age group per 100,000 population in age group.
9. Rates are based on the National Center for Health Statistics, estimates of the July 1, 2005 United States resident population from the Vintage 2005 postcensal series, prepared under a collaborative arrangement with the U.S. Census Bureau, Internet release date August 16, 2006.
10. Prior to 2005, Idaho BRFSS reported drinking status among all adults (binge drinking = 5 or more drinks in single occasion). Beginning in 2005, Idaho reports gender specific status (5 or more for males, 4 or more for females, in single occasion).

**Public Health District Health Indicators
2005 Data**

Item	Idaho	District 1	District 2	District 3	District 4	District 5	District 6	District 7
Maternal/Child								
1) % unintended pregnancies	34.8	40.5	33.1	35.0	31.7	41.4	36.7	30.7
2a) % live births with adequate prenatal care in 2005 ¹	41.9	44.4	47.5	36.3	42.5	45.8	39.1	42.5
2b) % live births with intensive prenatal care in 2005 ¹	32.3	34.9	30.5	29.0	35.6	28.0	38.2	27.7
2c) % live births with adequate or intensive prenatal care in 2005 ¹	74.2	79.3	78.0	65.3	78.1	73.8	77.3	70.2
3a) % live births with adequate prenatal care in 2004 ¹	41.6	45.6	48.2	37.2	39.9	47.2	39.1	41.6
3b) % live births with intensive prenatal care in 2004 ¹	32.6	29.0	29.1	30.6	39.9	27.5	39.1	25.4
3c) % live births with adequate or intensive prenatal care in 2004 ¹	74.2	74.6	77.3	67.8	79.8	74.7	78.2	67.0
4) % live births with low birth weight ²	6.7	6.5	5.7	6.6	6.9	6.5	7.6	6.3
5) % live births with tobacco use during pregnancy ³	13.2	22.5	16.0	12.3	12.3	13.7	13.5	8.4
6) % WIC participation ⁴	38.5	40.1	38.4	44.6	22.9	44.0	51.9	42.9
7) % prevalence of breastfeeding ⁵	88.0	88.4	89.9	90.0	90.1	84.1	83.7	88.1
Adolescents								
1) Teen pregnancy rate (15-19) ⁶	44.7	42.4	27.2	65.6	44.0	59.1	42.8	27.8
2) Motor vehicle crash death rate, 2003-2005 (ages 15-19) ^{7,9}	28.2	32.1	36.3	15.5	11.3	52.0	42.0	30.5
3) Suicide rate, 2003-2005 (ages 10-18) ^{8,9}	5.2	5.1	5.6	5.3	5.5	1.4	4.3	9.0

Item	Idaho	District 1	District 2	District 3	District 4	District 5	District 6	District 7
Adults								
1) % without health care coverage	18.9	21.2	22.5	25.7	12.9	24.7	15.1	16.3
2) % with no leisure time activity	21.6	20.8	23.9	27.7	17.3	24.4	21.0	21.1
3) % overweight (BMI \geq 25)	61.4	59.4	61.1	68.0	58.3	62.1	63.5	59.7
4) % diagnosed with diabetes	6.8	6.3	6.9	8.4	6.0	7.3	7.1	6.7
5) % who smoke	17.9	20.7	19.3	21.8	16.2	20.5	15.9	12.1
6) % who binge drink ¹⁰	14.2	13.2	13.6	13.3	19.6	12.5	10.4	7.5
7) % of females without breast cancer screening	na	na	na	na	na	na	na	na
8) % of males without prostate cancer screening	na	na	na	na	na	na	na	na
9) % who did not wear seatbelts	23.8	17.5	27.3	21.4	19.0	25.0	32.5	34.0
10) Suicide rate, 2003-2005 (ages 65+) ^{8,9}	22.9	26.6	33.3	26.4	15.0	19.8	22.6	23.6
11) % with no dental visit in the past 12 months	33.5	38.3	37.3	39.9	26.2	39.5	32.1	29.6

"na" indicates data are not available for this year.

SOURCES:

For Maternal/Child indicator 1: Pregnancy Risk Assessment Tracking Systems (PRATS), Bureau of Health Policy and Vital Statistics. PRATS is a representative sample of resident women aged 18+ who gave birth in Idaho.

For Maternal/Child indicators 2-7, Adolescents indicators, and Adults indicator 10: Vital Statistics, Bureau of Health Policy and Vital Statistics.

For Adults indicators 1-9, 11: Idaho Behavioral Risk Factor Surveillance System (BRFSS), Bureau of Health Policy and Vital Statistics.

NOTES:

1. In 2004, the Idaho birth certificate was revised. Beginning in 2004, Idaho prenatal care data are based on date of first prenatal care visit as reported in the mother's medical record. 2004 and 2005 APNCU data are not comparable with APNCU data based on month prenatal care began. The APNCU index classifies care as intensive, adequate, intermediate, inadequate, or no care by comparing the number of actual prenatal care visits that a woman was expected to receive, given the onset of prenatal care and the length of gestation. Percents based on records with stated data for APNCU calculation (date of first prenatal care visit, number of visits, and length of gestation). Data for 2004 have been corrected from the 2004 Health Indicators; 2004 data are comparable to 2005 data.
2. Percents based on records with stated birth weight.
3. Prior to 2004, tobacco use during pregnancy (yes or no) was reported on the Idaho birth certificate. Beginning in 2004, cigarette smoking before and during pregnancy is reported on the revised birth certificate. Data in 2004 and 2005 are not comparable to data prior to 2004.
4. The revised 2004 Idaho birth certificate includes a new variable for reporting if mother received WIC food for herself during this pregnancy (yes or no)?
5. The revised 2004 birth certificate includes a new variable for reporting if infant was being breastfed at birth or at the time the birth certificate was completed (yes or no)?
6. Pregnancies include live births, induced abortions, and reportable stillbirths (only those fetal deaths with a gestational period of 20+ weeks or which weight 350+ grams are required to be reportable under Idaho law.) Rate: Number of pregnancies per 1,000 females in corresponding age group. Rates are based on July 1, 2005 female population estimates from the National Center for Health Statistics, United States resident population from the Vintage 2005 postcensal series, prepared under a collaborative arrangement with the U.S. Census Bureau, Internet release date August 16, 2006.
7. Motor Vehicle Crash Death Rate: number of motor vehicle crash deaths from 2003 to 2005 per 100,000 population.
8. Suicide rate: number of suicides from 2003 to 2005 per 100,000 population.
9. Rates are based on the National Center for Health Statistics, estimates of the July 1, 2004 United States resident population from the Vintage 2004 postcensal series, prepared under a collaborative arrangement with the U.S. Census Bureau, Internet release date September 9, 2005.
10. Prior to 2005, Idaho BRFSS reported drinking status among all adults (binge drinking = 5 or more drinks in single occasion). Beginning in 2005, Idaho reports gender specific status (5 or more for males, 4 or more for females, in single occasion).